

CERTIFICATE OF ANALYSIS

Product BIOTINYLATED ANTI-MOUSE IgG (H+L)

AFFINITY PURIFIED

Catalog BA-2000

Amount <u>1.5 mg biotinylated IgG</u>

Concentration 1.5 mg/ml

Lot No. ZF1127

Produced in <u>horse</u>

Solution 10 mM phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide. Immunohistochemical

grade bovine serum albumin has been added for stabilization.*

Storage Refrigerate; for long-term storage, aliquots may be stored frozen.

Immunoglobulin biotinylated 99%

Antibody Activity after biotinylation (by solid phase IgG binding assay) 93%

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u>	Percent Cross-reactivity
hamster IgG, mouse IgM, rat IgG	12%
dog IgG, guinea pig IgG, swine IgG	3%
rat IgM, Rhesus Monkey IgG	1%
bovine IgG, bovine IgM, cat IgG, chicken IgG, dog IgM, goat IgG, goat IgM, human IgG, human IgM, rabbit IgG, rabbit IgM, sheep IgG,	<1%
sheep IgM, swine IgM	

If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

Vector Laboratories, Inc., 30 Ingold Road, Burlingame, CA 94010 U.S.A. Tel: (650)697-3600 • Fax: (650)697-0339 • Email: vector@vectorlabs.com • Web: www.vectorlabs.com

^{*} This product is at the same concentration as that provided in the VECTASTAIN $^{\text{®}}$ ABC kits. The recommended concentration range for use is 2-10 μ g/ml.