

## CERTIFICATE OF ANALYSIS

Product <u>BIOTINYLATED ANTI-MOUSE IgG (H+L)</u>

\*\*AFFINITY PURIFIED\*\* \*RAT ADSORBED\*

Catalog No. <u>BA-2001</u>

Amount <u>0.5 mg biotinylated IgG</u>

Concentration <u>0.5 mg/ml</u>

Lot No. <u>ZG0813</u>

Produced in <u>horse</u>

Solution 10 mM phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide.

Immunohistochemical grade bovine serum albumin has been added for

stabilization.

Storage Refrigerate; for long-term storage, aliquots may be stored frozen.

Immunoglobulin biotinylated 100%

Antibody Activity after biotinylation (by solid phase IgG binding assay) 90%

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u>	Percent Cross-reactivity
mouse IgM	25%
rat IgG	1.6%
dog IgG, guinea pig IgG, rat IgM, swine IgG	1%
bovine IgG, bovine IgM, cat IgG, chicken IgG, dog IgM, goat IgG, goat IgM, hamster IgG, human IgG, human IgM,	
rabbit IgG, rabbit IgM, Rhesus monkey IgG,	<1%
sheep IgG, sheep IgM, swine IgM	

If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

The recommended concentration range for use is  $2-10 \mu g/ml$ .