

CERTIFICATE OF ANALYSIS

Product BIOTINYLATED ANTI-HUMAN IgG (H+L)

AFFINITY PURIFIED

Catalog No. BA-3000

Amount 1.5 mg biotinylated IgG

Concentration <u>1.5 mg/ml</u>

Lot No. ZE0614

Produced in goat

Solution 10 mM phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide.

Immunohistochemical grade bovine serum albumin has been added for

stabilization.*

Storage Refrigerate; for long-term storage, aliquots may be stored frozen.

Immunoglobulin biotinylated 99%

Antibody Activity after biotinylation (by solid phase IgG binding assay) 95%

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u>	Percent Cross-reactivity
Rhesus monkey IgG human IgM	25% 12%
horse IgG, rat IgM, swine IgM	6%
cat IgG, dog IgG, dog IgM, guinea pig IgG, hamster IgG, mouse IgM, swine IgG	3%
bovine IgG, bovine IgM, chicken IgG, mouse IgG, rabbit IgG, rabbit IgM, rat IgG, sheep IgG, sheep IgM	<1%

If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

^{*} This product is at the same concentration as that provided in the VECTASTAIN $^{\circ}$ ABC kits. The recommended concentration range for use is 2-10 μ g/ml.