

CERTIFICATE OF ANALYSIS

Product <u>BIOTINYLATED ANTI-RAT IgG (H+L)</u>

AFFINITY PURIFIED

Catalog No. BA-4000

Amount <u>1.5 mg biotinylated IgG</u>

Concentration 1.5 mg/ml

Lot No. <u>ZF0305</u>

Produced in rabbit

Solution 10 mM phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide.

Immunohistochemical grade bovine serum albumin has been added for stabilization.

*

Storage Refrigerate; for long-term storage, aliquots may be stored frozen.

Immunoglobulin biotinylated 99%

Antibody Activity after biotinylation (by solid phase IgG binding assay) 93%

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u>	Percent Cross-reactivity
hamster IgG	25%
mouse IgM, rat IgM	12%
bovine IgG, horse IgG, mouse IgG	6%
guinea pig IgG, swine IgG	3%
bovine IgM, cat IgG, dog IgG, dog IgM, goat IgG,	
sheep IgG	1%
chicken IgG, goat IgM, human IgG, human IgM, sheep IgM, swine IgM	<1%

If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

Vector Laboratories, Inc., 30 Ingold Road, Burlingame, CA 94010 U.S.A.

^{*} This product has the same concentration as that provided in the VECTASTAIN [®] ABC kits. The recommended concentration range for use is 2-10 µg/ml.