

## CERTIFICATE OF ANALYSIS

Product BIOTINYLATED ANTI-RAT IgG (H+L)

\*\*AFFINITY PURIFIED\*\* \*MOUSE ADSORBED\*

Catalog <u>BA-9401</u>

Amount 0.5 mg active biotinylated IgG

Concentration <u>0.5 mg/ml</u>

Lot No. <u>ZG0608</u>

Produced in goat

Solution 10 mM phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide.

Immunohistochemical grade bovine serum albumin has been added for

stabilization.

Storage Refrigerate; for long-term storage, aliquots may be stored frozen.

Immunoglobulin biotinylated 100%

Antibody activity after biotinylation 95%

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u> <u>Percent Cross-reactivity</u>

hamster IgG 50%

rat IgM

bovine IgG, bovine IgM, cat IgG, chicken IgG, dog IgG, dog IgM, guinea pig IgG, horse IgG, human IgG, human IgM, mouse IgG, mouse IgM, rabbit IgG, rabbit IgM,

Rhesus monkey IgG, sheep IgG, sheep IgM, swine IgG.

<1%

Rhesus monkey IgG, sheep IgG, sheep IgM, swine IgG,

swine IgM

If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

The recommended concentration range for use is 2-10  $\mu$ g/ml.