

CERTIFICATE OF ANALYSIS

Product <u>BIOTINYLATED LECTIN KIT I</u>

Catalog No. <u>BK-1000</u>

Kit Lot ZG0612

Storage <u>2-8 °C</u>

Biotinylated Lectin	<u>Amount</u>	<u>Lot</u>	<u>Active</u> <u>Conjugate</u>	<u>Lectin</u> Biotinylated
Con A	1 mg	ZG0413	99%	99%
SBA	1 mg	ZG0612	99%	100%
WGA	1 mg	ZG0413	98%	100%
DBA	1 mg	ZG0413	93%	99%
UEA I	1 mg	ZG0413	98%	97%
RCA_{120}	1 mg†	ZG0413	99%	100%
PNA	1 mg	ZG0413	98%	99%

Solution 10 mM HEPES, 0.15 M NaC1, 0.1 mM Ca++, 0.08% sodium azide, pH 7.5.

Biotinylated Con A and PNA also contain 0.01 mM Mn++.

Concentration 2mg/ml

†Biotinylated *Ricinus communis* Agglutinin I (RCA₁₂₀) is at a concentration of 2.0 mg/ml in 10 mM phosphate, 0.15 M NaCl, 0.08% sodium azide, pH 7.5. **EXERCISE CAUTION IN ITS USE!** It has been found to have moderate toxicity in some animal models, but its toxicity to humans has not been fully determined, and a related protein, RCA₆₀, is highly toxic to humans. Never pipet solutions of this lectin by mouth. **DO NOT LYOPHILIZE!**

The recommended concentration range for use is 5-20 µg/ml.

Abbreviations:

CON A: Concanavalin A

SBA: Glycine max (soybean) agglutinin

WGA: Triticum vulgaris (wheat germ) agglutinin

DBA: Dolichos biflorus agglutinin