

CERTIFICATE OF ANALYSIS

Product <u>DyLight®594 ANTI-GOAT IgG (H+L)</u>

AFFINITY PURIFIED

Catalog No. DI-3094

Amount <u>1.5 mg active conjugate</u>

Lot No. ZF0415

Produced in <u>horse</u>

Concentration 1.5 mg/ml active conjugate

Solution 10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide.

Maximum Excitation @ 592 nm

Maximum Emission @ 617 nm

F/P (molar) 4.1


Storage Conditions Refrigerate in the dark.


Recommended concentration range for use 5-20 μg/ml

The affinity purified antibody has been extensively adsorbed on solid phases and solid-phase enzyme immunoassay shows minimal cross-reactivity against the following immunoglobulins*:

human, mouse, rabbit, pig, cat, dog, chicken, monkey, guinea pig, hamster, rat.

*The use of bovine products such as casein, serum, albumin or non-fat dry milk as blocking agents may produce a high background due to cross-reactivity of anti-goat IgG with bovine immunoglobulins that may be present. Blocking with Normal Horse Serum (Cat. No. S-2000) or Animal-Free Blocker[™] solution (Cat. No. SP-5030) is recommended.





DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.