



CERTIFICATE OF ANALYSIS

Product **GLYCOPROTEIN ELUTING SOLUTION**
for SIALIC ACID-BINDING LECTINS

Catalog No. ES-7100

Amount 100 ml

Lot No. ZF0506

Storage Room Temperature

Test Results

This solution elutes >90% of Sialic Acid-labeled BSA (Cat. No. G-7000) from *Sambucus Nigra* Lectin Agarose (Cat. No. AL-1303). *

Protocol

- 1) Wash lectin agarose column to remove unbound proteins with HEPES- or TRIS-Buffered saline (HBS, TBS), pH 7.5, until OD280nm reaches background level.
- 2) Apply Glycoprotein Eluting Solution and allow to flow through the column by gravity. Note: Do not apply pressure to speed elution because this may reduce the percent recovery of the bound glycoprotein. Complete elution may take up to 10 column volumes.*
- 3) Following elution, the agarose lectin column is ready for reuse after equilibrating with 2-4 column volumes of HBS or TBS, pH 7.5.

*These results may not be typical for other glycoproteins, since oligosaccharide structures of glycoproteins vary significantly. The eluting solution has a high osmolarity. If downstream applications are adversely affected by salt, eluted glycoproteins may require dialysis or gel filtration.