

CERTIFICATE OF ANALYSIS

Product PEROXIDASE ANTI-MOUSE IgG (H+L)
AFFINITY PURIFIED

Catalog No. PI-2000

Amount <u>1 mg active conjugate</u>

Lot No. <u>ZF0503</u>

Produced in <u>horse</u>

Concentration <u>1 mg/ml active conjugate</u>

Solution 10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.05% thimerosal

Molar ratio of active conjugate Moles of

 $\frac{\text{Moles of HRP}}{\text{Moles of A + 3.3}} = 3.3$

Moles of Antibody

Storage Conditions Refrigerate; DO NOT FREEZE

Remarks: Do not use sodium azide with this product since it is an inhibitor of horseradish peroxidase.

Solid-phase enzyme immunoassay of cross-reactivity:

Protein	Percent Cross-reactivity
mouse IgM	25%
hamster IgG, rat IgG	6%
bovine IgG, dog IgG, guinea pig IgG, Rhesus monkey IgG, swine IgG	3%
dog IgM, swine IgM	1%
bovine IgM, cat IgG, chicken IgG, goat IgG, goat IgM, guinea pig IgM, human IgG, human IgM, rabbit IgG, rabbit IgM, rat IgM, sheep IgG,	:10/
sheep IgM	<1%

If this peroxidase-labeled antibody is to be used in tissues which may contain cross-reacting endogenous immunoglobulins, dilution of this antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

Recommended diluent for microtitration plate assays: 10 mM phosphate, 0.15 M NaCl, pH 7.5, 0.1% Tween 20. Working concentration of HRP-anti mouse IgG is 0.5-5.0 µg/ml.

Vector Laboratories, Inc., 30 Ingold Road, Burlingame, CA 94010 U.S.A.

Tel: (650)697-3600 • Fax: (650)697-0339 • Email: vector@vectorlabs.com • Web: www.vectorlabs.com