

CERTIFICATE OF ANALYSIS

Product PEROXIDASE ANTI-HUMAN IgG (H+L)
AFFINITY PURIFIED

Catalog No. PI-3000

Amount <u>1 mg active conjugate</u>

Lot No. ZE1129

Produced in goat

Concentration <u>1 mg/ml active conjugate</u>

Solution 10 mM HEPES, 0.15 M NaCl, pH 7.5, preservative

Molar ratio of active conjugate Moles of HRP

Moles of Antibody

Storage Conditions Refrigerate; DO NOT FREEZE

Remarks: Do not use sodium azide with this product since it is an inhibitor of horseradish peroxidase.

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u>	Percent Cross-reactivity
guinea pig IgG, Rhesus monkey IgG	25%
cat IgG, dog IgM, human IgM	6%
chicken IgG, dog IgG, horse IgG, swine IgG	3%
hamster IgG, mouse IgM	1%
bovine IgG, bovine IgM, mouse IgG, rabbit IgG	
rat IgG, sheep IgG	<1%

If this peroxidase-labeled antibody is to be used in tissues which may contain cross-reacting endogenous immunoglobulins, dilution of this antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

Recommended diluent for microtitration plate assays: 10 mM phosphate, 0.15 M NaCl, pH 7.5, 0.1% Tween 20. Working concentration of HRP-anti human IgG is 0.5-5.0 μg/ml.

Vector Laboratories, Inc., 30 Ingold Road, Burlingame, CA 94010 U.S.A.

Tel: (650)697-3600 • Fax: (650)697-0339 • Email: vector@vectorlabs.com • Web: www.vectorlabs.com