

CERTIFICATE OF ANALYSIS

Product TEXAS RED® ANTI-MOUSE IgG (H+L)

AFFINITY PURIFIED

Catalog No. TI-2000

Amount 1.5 mg active conjugate

Lot No. ZF0829

Produced in horse

Concentration 1.5 mg/ml active conjugate

Solution 10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide.

Free-Fluorochrome None

F/P (molar) 2.5

Maximum Excitation @ 595-604 nm

Maximum Emission @ 606-615 nm

Storage Conditions Refrigerate in the dark.

Antibody Activity (by solid phase IgG binding assay) 95%

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u>	Percent Cross-reactivity
mouse IgM	25%
hamster IgG, rat IgG	6%
bovine IgG, dog IgG, guinea pig IgG, Rhesus monkey IgG, swine IgG	3%
dog IgM, rat IgM, swine IgM	1%
bovine IgM, cat IgG, chicken IgG, goat IgG, goat IgM, guinea pig	
IgM,	
human IgG, human IgM, rabbit IgG, rabbit IgM, sheep IgG,	<1%
sheep IgM	

If this fluorochrome antibody is to be used in tissues which may contain cross-reacting endogenous immunoglobulins, dilution of this antibody may be made in buffers containing 2% normal serum from the same species as the tissue. The recommended concentration range for use is 5-20 µg/ml.

Vector Laboratories, Inc., 30 Ingold Road, Burlingame, CA 94010 U.S.A.

^{*}J. Immunological Methods <u>50</u>, 193 (1982)

[®] Trademark of Molecular Probes