



Goat Anti-Biotin, Unconjugated SP-3000-1

Product Images



Short Description

Our high affinity polyclonal anti-biotin antibody is produced by hyperimmunizing goats with biotin and is purified by affinity chromatography. The unconjugated antibody can be detected with a VECTASTAIN® ABC system or the ImmPRESS® Anti-Goat Ig polymer detection system (MP-7405) for blot, immunohistochemical and in situ hybridization applications.

Additional Information

Unit Size	1 mg
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs
Recommended Storage	2-8 °C
Recommended Usage	Reconstitute by adding 1 ml water. The resulting solution will have the following composition: 10 mM sodium phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide, 20 mg/ml sucrose. This product is recommended for use in transfer blot applications (northern, Southern, western, dot blots, etc.), immunohistochemical, and in situ hybridization applications. The recommended concentration range for use is 0.5-5 µg/ml. This reagent can be detected with a VECTASTAIN® ABC System or the ImmPRESS™ Anti-Goat IgG Polymer Detection System (Cat. No. MP-7405).
Detection Target	Biotin
Conjugate	Unconjugated
Host Species	Goat
Format	Lyophilized

