



Goat Anti-Human IgM Antibody, Mu Chain Specific, Biotinylated

BA-3020-.5

[Product Images](#)



Short Description

Biotinylated Goat Anti-Human IgM Antibody, mu chain specific, is used in an avidin-biotin or streptavidin-biotin detection system. These antibodies can be used for tissue and cell staining, ELISAs, and blots. This mu chain-specific antibody is produced specifically to distinguish between chains or classes of target immunoglobulins.

Features:

- This mu chain specific antibody has virtually no cross-reactivity with other immunoglobulin classes or other heavy or light chains
- Included in the VECTASTAIN® ABC Kits
- Supplied in solution

Additional Information

Unit Size	0.5 mg
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs
Concentration	0.5 mg active conjugate/ml
Recommended Storage	2-8 °C; Store frozen for long term storage
Solution	10 mM sodium phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide, 1 mg/ml bovine serum albumin.
Recommended Usage	The recommended concentration range for use is 2-10 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue. This is the same solution as that provided in the VECTASTAIN® ABC kits.
Target Species	Human
Conjugate	Biotinylated
Host Species	Goat
Format	Concentrate

