



Goat Anti-Mouse IgG Antibody (H+L), Biotinylated

BA-9200-1.5

[Product Images](#)



Short Description

The biotinylated goat anti-mouse IgG secondary antibody is used in an avidin-biotin or streptavidin-biotin detection system. These antibodies can be used for tissue and cell staining, ELISAs, and blots.

Features:

- Recognizes both heavy and light chains (H+L)
- Supplied in solution

Additional Information

Unit Size	1.5 mg
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs
Concentration	1.5 mg active conjugate/ml
Recommended Storage	2-8 °C; Store frozen for long term storage
Solution	10 mM sodium phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide, 3 mg/ml bovine serum albumin.
Recommended Usage	The recommended concentration range for use is 2-10 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue. NOTE: This product is raised in goat and is not the same as that contained in the VECTASTAIN ABC Kits for the detection of mouse IgG which is raised in horse.
Target Species	Mouse
Conjugate	Biotinylated
Host Species	Goat
Format	Concentrate

