



Goat Anti-Peanut Agglutinin (PNA), Biotinylated

BA-0074-.5

[Product Images](#)



Short Description

These antibodies to lectins are produced by immunizing goats with pure lectins. Following conventional purification steps, specific antibody is isolated by affinity chromatography on lectin-agarose columns. Each anti-lectin is supplied lyophilized in buffered saline.

Biotinylated anti-lectins are optimally labeled with biotin.

Additional Information

Unit Size	0.5 mg
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, Blotting Applications, Elispot, ELISAs, Glycobiology
Concentration	1 mg active conjugate/ml
Recommended Storage	2-8 °C; Store frozen for long term storage
Solution	10 mM sodium phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide, 20 mg/ml sucrose
Recommended Usage	The recommended concentration range for use is 5-20 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.
Detection Target	Lectin
Conjugate	Biotinylated
Host Species	Goat
Format	Concentrate

