



Goat Anti-Rabbit IgG Antibody (H+L), Biotinylated

BA-1000-1.5

[Product Images](#)



Short Description

The biotinylated goat anti-rabbit IgG secondary antibody is used in an avidin-biotin or streptavidin-biotin detection system. These antibodies can be used for tissue and cell staining, ELISAs, and blots.

Features:

- Recognizes both heavy and light chains (H+L)
- Cross-reactivities that are likely to interfere with specific labeling are removed by solid phase adsorption techniques
- Included in the VECTASTAIN® ABC kits

Additional Information

| | |
|---------------------|---|
| Unit Size | 1.5 mg |
| Applications | Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs |
| Concentration | 1.5 mg active conjugate/ml |
| Recommended Storage | 2-8 °C; Store frozen for long term storage |
| Solution | 10 mM sodium phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide, 3 mg/ml bovine serum albumin. |
| Recommended Usage | The recommended concentration range for use is 2-10 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue. Note: This product is at the same concentration as that provided in the VECTASTAIN® ABC kits. |
| Target Species | Rabbit |
| Conjugate | Biotinylated |
| Host Species | Goat |
| Format | Concentrate |

