



---

# Goat Anti-Rabbit IgG Antibody (H+L), Peroxidase

## PI-1000-1

[Product Images](#)

---



## Short Description

---

The HRP Goat Anti-Rabbit IgG Antibody binds to rabbit primary antibodies for tissue staining, ELISAs, and blots. It can be detected using a horseradish peroxidase substrate.

### Features:

- Recognizes both heavy and light chains (H+L)
- Cross-linked with the highest specific activity horseradish peroxidase
- Supplied in solution

## Additional Information

---

Unit Size	1 mg
Applications	Immunohistochemistry / Immunocytochemistry, In situ hybridization, Blotting Applications, Elispot, ELISAs
Concentration	1 mg active conjugate/ml
Recommended Storage	2-8 °C, DO NOT FREEZE
Solution	10 mM HEPES, 0.15 M NaCl, pH 7.5, preservative
Recommended Usage	Recommended diluent for microtitration plate assays: 10 mM phosphate, 0.15 M NaCl, pH 7.5, 0.1% Tween 20. Working concentration of HRP-anti rabbit IgG is 0.5-5.0 µg/ml. Do not use sodium azide with this product since it is an inhibitor of horseradish peroxidase.
Target Species	Rabbit
Conjugate	Peroxidase
Host Species	Goat
Format	Concentrate
Safety Title	Cancer

