




# Goat Anti-Streptavidin Antibody, Biotinylated

## BA-0500-.5

[Product Images](#)



## Short Description

---

Biotinylated Anti-Streptavidin has been widely used as an amplifying reagent in immunohistochemistry, *in situ* hybridization, microarray assays, ELISAs, blots, and many other applications. The capability of binding streptavidin via either biotin binding sites or through antigen binding sites, makes this biotinylated antibody unique. Anti-Streptavidin does not bind avidin, and Anti-Avidin does not recognize streptavidin.

## Additional Information

---

Unit Size	0.5 mg
Applications	Immunofluorescence, In situ hybridization
Concentration	0.5 mg active conjugate/ml
Recommended Storage	2-8 °C; Store frozen for long term storage
Solution	10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.08% sodium azide, 20 mg/ml sucrose.
Recommended Usage	The recommended concentration range for use is 1-10 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.
Detection Target	Streptavidin
Conjugate	Biotinylated
Host Species	Goat
Format	Concentrate

