



Horse Anti-Goat IgG Antibody (H+L), Biotinylated, R.T.U.

BP-9500-50

[Product Images](#)



Short Description

The Ready-to-Use (R.T.U.) Biotinylated Horse Anti-Goat IgG Antibody is used in an avidin-biotin or streptavidin-biotin detection system. These antibodies can be used for tissue and cell staining, ELISAs, and blots.

Features:

- Suitable for use with goat, sheep, and bovine IgG primary antibodies
- Recognizes both heavy and light chains (H+L)

Additional Information

Unit Size	50 ml
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs
Concentration	Ready-to-use solution of biotinylated horse anti-goat IgG in PBS* containing carrier protein and 0.08% sodium azide as preservative.
Recommended Storage	2-8 °C
Recommended Usage	Following incubation with primary antibody, wash specimens in buffer. Apply RTU Biotinylated Anti-Goat IgG and incubate for 30 minutes at room temperature. Wash in buffer for 5 minutes and follow with avidin- or streptavidin-based detection reagent such as RTU VECTASTAIN® Elite® ABC Reagent (PK-7100). * 10mM sodium phosphate, pH 7.8, 0.15M NaCl.
Target Species	Goat
Conjugate	Biotinylated
Host Species	Horse
Format	Ready-to-Use

