



---

# Horse Anti-Goat IgG Antibody (H+L), Peroxidase

## PI-9500-1

[Product Images](#)

---



## Short Description

---

### Features:

- Affinity-purified, ultrapure, high affinity antibody
- Suitable for use with goat, sheep, and bovine IgG primary antibodies
- Thoroughly adsorbed against serum and immunoglobulins from potentially interfering species
- Unless otherwise specified, antibodies recognizes both heavy and light chains (H+L)
- Cross-linked with the highest specific activity horseradish peroxidase
- Supplied in solution

### Additional Information

---

Unit Size	1 mg
Applications	Immunohistochemistry / Immunocytochemistry, In situ hybridization, Blotting Applications, Elispot, ELISAs
Concentration	1 mg active conjugate/ml
Recommended Storage	2-8 °C, DO NOT FREEZE
Solution	10 mM HEPES, 0.15 M NaCl, pH 7.5, preservative
Recommended Usage	<p>Important Note: The use of bovine products such as casein, serum, albumin or non-fat dry milk as blocking agents in ELISA or blot assays may produce a high background due to cross-reactivity of anti-goat IgG with bovine immunoglobulins that may be present. If this antibody is to be used in these applications, blocking with Animal-Free blocker™ solution (Cat. No. SP-5030) is recommended. Recommended diluent for microtitration plate assays: 10 mM phosphate, 0.15 M NaCl, pH 7.5, 0.1% Tween 20. Working concentration of HRP-anti goat IgG is 0.5-5.0 µg/ml.</p>
Target Species	Goat
Conjugate	Peroxidase
Host Species	Horse
Format	Concentrate
Safety Title	Cancer

