



---

# Wisteria Floribunda Lectin (WFA, WFL), Agarose bound

## AL-1353-2

[Product Images](#)

---



## Short Description

---

### Features:

- Matrix is heat stable, cross-linked 4% agarose beads with a molecular exclusion of about  $2 \times 10^7$  daltons
- Bead diameter ranges in size from 45-165 microns
- Matrix is stable in solutions at pH 3-11 as well as many organic solvents
- Immobilized lectins are prepared using affinity purified lectins
- Covalent attachment preserves lectin activity and minimizes conformational changes that might result in nonspecific or hydrophobic interactions
- Hydrophilic spacer arm is inserted between the lectin and the matrix
- Conjugated proteins are not leached off the beads by Tris or other routinely used buffers
- No residual charges present after conjugation. This minimizes non-specific binding to the matrix
- Product supplied as a 1:1 suspension in buffer
- 3 mg lectin/ml gel
- Inhibiting/Eluting Sugar: 200 mM *N*-acetylgalactosamine or Glycoprotein Eluting Solution (ES-2100)

## Additional Information

---

Unit Size	2 ml
Applications	Glycobiology, Affinity Chromatography
Recommended Storage	2-8 °C DO NOT FREEZE
Solution	10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.1 mM CaCl <sub>2</sub> , 20 mM galactose, 0.08% sodium azide
Recommended Usage	Wash gel thoroughly with buffer before use to remove sugar added to stabilize the lectin. Recommended product for eluting glycoconjugates bound to this agarose-lectin: Glycoprotein Eluting solution, Cat. No. ES-2100. Alternatively, 200 mM <i>N</i> -acetylgalactosamine or several column volumes of 200 mM acetic acid can be used. After use, wash the gel with several column volumes of buffered saline, then resuspend gel in buffered saline containing 0.08% sodium azide for storage.
Matrix Conjugate	Lectins
Sugar Specificity	<i>N</i> -Acetylgalactosamine
Conjugate	Agarose

