



Wisteria Floribunda Lectin (WFA, WFL), Fluorescein

FL-1351-2

[Product Images](#)



Short Description

The binding specificity of *Wisteria floribunda* lectin (WFL) is not completely clear but this lectin appears to preferentially bind carbohydrate structures terminating in *N*-acetylgalactosamine linked α or β to the 3 or 6 position of galactose. This lectin has been used to fractionate lymphocyte populations, and although not mitogenic, elicits the production of lymphokines from murine splenocytes.

Fluorescein labeled WFL has an appropriate number of fluorochromes bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated fluorochromes.

- Excitation maximum: 495 nm
- Emission maximum: 515 nm
- Color: Green

Additional Information

Unit Size	2 mg
Applications	Immunofluorescence, Glycobiology
Recommended Usage	The recommended concentration range for use is 5-20 μ g/ml.
Recommended Storage	2-8°C
Maximum Excitation	495-500 nm
Inhibiting and/or Eluting Sugar	200 mM <i>N</i> -acetylgalactosamine(S-9001)
Maximum Emission	514-521 nm
Solution	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide. 0.1 mM CaCl ₂
Concentration	2 mg active conjugate/ml
Conjugate	Fluorescein
Color of Fluorescence	Green
Sugar Specificity	<i>N</i> -Acetylgalactosamine

