



# Agarose bound Concanavalin A (Con A)

## Product Images



## Short Description

---

Agarose bound Con A is prepared using our affinity-purified lectins. Con A recognizes  $\alpha$ -linked mannose present as part of a core oligosaccharide in many serum and membrane glycoproteins. At neutral and alkaline pH, Con A exists as a tetramer of four identical subunits; below pH 5.6, Con A dissociates into active dimers of 52 kDa. Acetylation, succinylation, or other derivatizations can also produce stable forms with dimeric structures. (See succinylated Con A). Nicks in the sequence are often present in the purest preparations due to hydrolytic damage within the seeds.

### Features:

- Bead diameter ranges in size from 45-165 microns
- Matrix is stable in solutions at pH 3-11 as well as many organic solvents
- Immobilized lectins are prepared using affinity purified lectins
- Covalent attachment preserves lectin activity and minimizes conformational changes that might result in nonspecific or hydrophobic interactions
- Hydrophilic spacer arm is inserted between the lectin and the matrix
- Conjugated proteins are not leached off the beads by Tris or other routinely used buffers
- No residual charges present after conjugation. This minimizes non-specific binding to the matrix
- Product supplied as a 1:1 suspension in buffer

## Additional Information

Unit Size	10 ml, 100 ml
Applications	Glycobiology, Affinity Chromatography
Recommended Storage	2-8 °C DO NOT FREEZE
Solution	10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.1 mM CaCl <sub>2</sub> , 0.01 mM MnCl <sub>2</sub> , 20 mM glucose, 0.08% sodium azide
Recommended Usage	Wash gel thoroughly with buffer before use to remove sugar added to stabilize the lectin. Recommended product for eluting glycoconjugates bound to this agarose-lectin: Glycoprotein Eluting Solution, Cat. No. ES-1100. Alternatively, a 0.2 M α-methyl mannoside/0.2 M α-methyl glucoside mixture can be used. After use, wash the gel with several column volumes of buffered saline, then resuspend gel in buffered saline containing 0.08% sodium azide for storage.
Matrix Conjugate	Lectins
Sugar Specificity	Mannose, Glucose
Conjugate	Agarose

### Agarose bound Concanavalin A (Con A)

#### Features:

- Matrix is heat stable, cross-linked 4% agarose beads with a molecular exclusion of about  $2 \times 10^7$  daltons
- Bead diameter ranges in size from 45-165 microns
- Matrix is stable in solutions at pH 3-11 as well as many organic solvents
- Immobilized lectins are prepared using affinity purified lectins
- Covalent attachment preserves lectin activity and minimizes conformational changes that might result in nonspecific or hydrophobic interactions
- Hydrophilic spacer arm is inserted between the lectin and the matrix
- Conjugated proteins are not leached off the beads by Tris or other routinely used buffers
- No residual charges present after conjugation. This minimizes non-specific binding to the matrix
- Product supplied as a 1:1 suspension in buffer
- 6 mg lectin/ml gel
- Inhibiting/Eluting Sugar: mixture of 200 mM  $\alpha$ -methylmannoside/200 mM  $\alpha$ -methylglucoside or Glycoprotein Eluting Solution (ES-1100)



## Agarose bound Concanavalin A (Con A)

### Features:

- Matrix is heat stable, cross-linked 4% agarose beads with a molecular exclusion of about  $2 \times 10^7$  daltons
- Bead diameter ranges in size from 45-165 microns
- Matrix is stable in solutions at pH 3-11 as well as many organic solvents
- Immobilized lectins are prepared using affinity purified lectins
- Covalent attachment preserves lectin activity and minimizes conformational changes that might result in nonspecific or hydrophobic interactions
- Hydrophilic spacer arm is inserted between the lectin and the matrix
- Conjugated proteins are not leached off the beads by Tris or other routinely used buffers
- No residual charges present after conjugation. This minimizes non-specific binding to the matrix
- Product supplied as a 1:1 suspension in buffer
- 6 mg lectin/ml gel
- Inhibiting/Eluting Sugar: mixture of 200 mM  $\alpha$ -methylmannoside/200 mM  $\alpha$ -methylglucoside or Glycoprotein Eluting Solution (ES-1100)

