




5(6)-TAMRA ACID

SKU: FP-1251



DESCRIPTION

488/532



Laser
line

TRITC



Common
filter set

556



Excitation
max

573



Emission
max

5(6)-TAMRA Acid (5(6)-Carboxytetramethylrhodamine, TMR, TRITC) is a bright orange-fluorescent dye with excitation ideally suited to the 532 nm or 546 nm laser lines. It has been

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



used widely for preparing peptide, protein, nucleotide and nucleic acid conjugates, especially fluorescent antibodies and avidin derivatives used in immunochemistry. The absorbance and emission maxima of TAMRA conjugates are 553 nm and 575 nm respectively.


TAMRA Acid is a reagent of choice for the preparation of custom activated esters that often are not commercially available. Examples of such activated esters include sulfo-NHS, TFP (2,3,5,6-Tetrafluorophenol), STP (4-Sulfo-2,3,5,6-Tetrafluorophenol, Sodium Salt). Another common application for non-activated carboxylic acid is peptide modification during solid phase synthesis, which usually requires in-situ activation with peptide coupling reagents, for example HATU. TAMRA Acid is also often used for control experiments, and for calibration.

SPECIFICATIONS

Molecular Weight	430.46
Extinction Coefficient	92,000 cm ⁻¹ M ⁻¹
Reactivity	Primary amines
Unit Size	25 mg, 100 mg, 1000 mg
Solubility	DMSO, DMF, MeOH
Storage Instructions	-20°C.
Spectrally Similar Dyes	Alexa Fluor® 546, TAMRA, CF™ 543, MB™ 543
Excitation/Emission Maximum	553/575 nm
Shipping Conditions	Ambient temperature
Shipping Instructions	Ambient temperature

ABS/EM SPECTRA


For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



DOCUMENTS

- [Safety Data Sheet](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.