




DADPS H2/D2 BIOTIN ALKYNE PACK - 2 VIALS

SKU: CCT-1451



DESCRIPTION

While there has been much interest in profiling the intact glycoproteome, the complexity of glycoproteoforms (and more broadly, all proteoforms) remains challenging to completely define. Mass spectrometry (MS) is commonly employed for characterization of complex proteomic samples. A popular strategy for protein identification is the bottom-up shotgun proteomics approach. In this method, a mixture of proteins is subjected to proteolytic digestion, the resulting peptides are separated by LC and detected by MS, and their parent proteins are inferred from the assigned peptide sequences.

To convert MS data acquired from proteolytic digests into protein identifications, tandem MS can be used to obtain sequence information for individual peptides, followed by comparing an in-silico proteolytic digest of an organism's proteome. Typically, only the most abundant peptides are selected for fragmentation (Figure 2), whereas data for those peptides in relatively low quantities are not obtained. An inherent problem in shotgun proteomics is identifying proteins of low abundance, such as biomarkers for disease states, against a background of proteins whose concentrations can span up to 12 orders of magnitude.

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.




Figure 1. Metabolic labeling with a chemically functionalized glycan, chemical tagging and enrichment using an isotopic recoding affinity probe To address the unique challenges of the global characterization of the intact glycoproteome, a mass-independent chemical glycoproteomics platform, termed *isotope targeted glycoproteomics* (IsoTag) was developed by the Carolyn Bertozzi group. The platform is comprised of four central components: (i) metabolic labeling with a chemically functionalized glycan, (ii) chemical tagging and enrichment using an isotopic recoding affinity probe, (iii) directed tandem MS, and (iv) targeted glycopeptide assignment (Figure 2).




Figure 2. Traditional proteomics and Iso-Tag-directed proteomics workflow IsoTag is performed by isotopic recoding and enrichment of metabolically labeled glycoproteins followed by directed tandem MS (MS2 or MSn) analysis and intact glycopeptide assignment. Isotopic recoding is

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



accomplished by metabolic labeling of cell or tissue samples with azide- or alkyne-functionalized sugars, followed by chemical conjugation with a biotin probe bearing a unique isotopic signature.

Some examples of sugar labels are peracetylated N-azidoacetylmannosamine (Ac4ManNAz), which is converted to the corresponding azidosialic acid (SiaNAz), and peracetylated N-azidoacetylgalactosamine (Ac4GalNAz), which is metabolized to label glycans possessing N-acetylglucosamine (GlcNAc) or N-acetylgalactosamine (GalNAc) (not provided with kit).

In order to perform isotopic tagging, the kit provides two cleavable IsoTaG probes encoded by zero [M] and two [M + 2] deuterium atoms. Probes with different encoding can be provided by Click Chemistry Tools through custom synthesis. The IsoTaG probe with zero, and that with two deuterium atoms [M, M + 2] can be used in different proportions; 1:1, 1:2, 1:3 and 1:4. Pattern recognition with isotopic ratio of 1:3 showed the highest fidelity.




Figure 3. Cleavable IsoTaG probe encoded by zero deuterium atoms [M] (R = H) and two deuterium atoms [M+2] (R = D). Through these probes, a unique isotopic signature is embedded exclusively into the glycopeptides. The isotopic signature serves as a computationally recognizable full-scan MS reporter. A computational algorithm, termed isotopic signature transfer and mass pattern prediction (IsoStamp), for the detection of recorded species in full-scan mass spectra, was also developed by the Carolyn Bertozzi group. IsoStamp compares observed and predicted isotopic envelopes to identify chemically tagged species in full-scan mass spectra.

IsoTag has the potential to enhance any proteomics platform that employs chemical labeling for targeted protein identification, including isotope-coded affinity tagging, isobaric tagging for relative and absolute quantitation, and chemical tagging strategies for post-translational modification.

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



SPECIFICATIONS

Molecular Weight	844.41
Appearance	Oil to grey amorphous solid
Unit Size	3 vials
Solubility	DMSO, DMF
Storage Instructions	-20C
DADPS D2 Biotin Azide	1 vial, 2 mg
DADPS H2 Biotin Azide	1 vial, 2 mg
Purchase Note	Add some note
Shelf Life	3 years at -20C
Shipping Instructions	Frozen

SELECTED REFERENCES

1. Woo, C.M., *et al.* (2017). Development of IsoTaG, a Chemical Glycoproteomics Technique for Profiling Intact N- and O-Glycopeptides from Whole Cell Proteomes. *J. Proteome Res.*, **16**: 1706–18. [[PubMed](#)]
2. Woo, C.M., *et al.* (2017). Mapping and Quantification of Over 2000 O-linked Glycopeptides in Activated Human T Cells with Isotope-Targeted Glycoproteomics (Isotag). *Mol. Cell. Proteomics* **17**: 764-75. [[PubMed](#)]
3. Goa, G., *et al.* (2017). Small Molecule Interactome Mapping by Photoaffinity Labeling Reveals Binding Site Hotspots for the NSAIDs. *J. Am. Chem. Soc.*, **140**: 4259–68. [[PubMed](#)]
4. Woo, C.M., *et al.* (2015). Isotope-targeted glycoproteomics (IsoTaG): a mass-independent platform for intact N- and O-glycopeptide discovery and analysis. *Nat Methods.*, **12**: 561-7. [[PubMed](#)]
5. Weerapana, E., *et al.* (2010). Quantitative reactivity profiling predicts functional cysteines in proteomes. *Nature* **468**: 790–5. [[PubMed](#)]

DOCUMENTS

- [Safety Data Sheet](#)
- [Datasheet](#)

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.




www.vectorlabs.com

Email: customerservice@vectorlabs.com

Telephone: [\(650\) 697-3600](tel:(650)697-3600)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.

[DADPS H2/D2 Biotin Alkyne Pack - 2 vials](https://vectorlabs.com/products/dadps-h2-d2-biotin-alkyne-pack/)

<https://vectorlabs.com/products/dadps-h2-d2-biotin-alkyne-pack/>