




AMCA ACID

SKU: FP-1235



DESCRIPTION

350



Laser
line

DAPI



Common
filter set

346



Excitation
max

442



Emission
max

AMCA Acid (7-amino-4-methylcoumarin-3-acetic acid) is one of the most popular blue fluorescent tagging molecules. Often used as contrasting probes for double- and triple-labeling immunofluorescence microscopy, arrays and in situ hybridization. The desirable properties of AMCA dyes include a relatively large Stoke's shift and resistance to photobleaching.

AMCA Acid is a reagent of choice for the preparation of custom activated esters that often are not commercially available. Examples of such activated esters include sulfo-NHS, TFP (2,3,5,6-Tetrafluorophenol), STP (4-Sulfo-2,3,5,6-Tetrafluorophenol, Sodium Salt). Another common application for non-activated carboxylic acid is peptide modification during solid phase synthesis, which usually requires in-situ activation with peptide coupling reagents, for example HATU. AMCA Acid is also often used for control experiments, and for calibration.


For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



SPECIFICATIONS

Molecular Weight	233.27
Extinction Coefficient	19,000 cm ⁻¹ M ⁻¹
Reactivity	Primary amines (needs activation)
Unit Size	25 mg, 100 mg, 1000 mg
Solubility	DMSO, DMF
Storage Instructions	-20°C.
Spectrally Similar Dyes	Alexa Fluor® 350, AMCA, DyLight® 350
Excitation/Emission Maximum	345/450 nm
Shipping Conditions	Ambient temperature
Shipping Instructions	Ambient temperature

ABS/EM SPECTRA



DOCUMENTS


- [Safety Data Sheet](#)

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.