



# GOAT ANTI-STREPTAVIDIN ANTIBODY, BIOTINYLATED

**SKU:** BA-0500-.5



---

## DESCRIPTION

Biotinylated Anti-Streptavidin has been widely used as an amplifying reagent in immunohistochemistry, *in situ* hybridization, microarray assays, ELISAs, blots, and many other applications. The capability of binding streptavidin via either biotin binding sites or through antigen binding sites, makes this biotinylated antibody unique. Anti-Streptavidin does not bind avidin, and Anti-Avidin does not recognize streptavidin.

## SPECIFICATIONS

|                             |                                                                         |
|-----------------------------|-------------------------------------------------------------------------|
| <b>Format</b>               | Concentrate                                                             |
| <b>Formulation</b>          | 10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.08% sodium azide, 20 mg/ml sucrose. |
| <b>Unit Size</b>            | 0.5 mg                                                                  |
| <b>Storage Instructions</b> | 2-8 °C; Store frozen for long term storage                              |

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



## Usage Summary

The recommended concentration range for use is 1-10 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

## Applications

Immunofluorescence, In situ hybridization

## Concentration

0.5 mg active conjugate/ml

## Conjugate

Biotinylated

## Reactive Species

Goat

## Host Species


Goat

# TECHNICAL INFORMATION

These antibodies can be used either as part of preformed complexes or in sequence to amplify fluorescent signals. When used in sequence, the target is first labeled with fluorochrome-conjugated avidin or streptavidin, followed by incubation with Biotinylated Anti-Avidin or Biotinylated Anti-Streptavidin, followed by a second layer of fluorochrome-conjugated avidin or streptavidin. This sequence can be repeated. This multi-layered approach introduces more fluorochromes at the target site and can provide a multi-fold amplification over a single layer.

These affinity purified antibodies are also available unconjugated or fluorescein-labeled.

The following figures illustrate different experimental setups used to achieve increasing levels of sensitivity.



**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



## CITATIONS




Powered by Bioz © 2023 See more details on Bioz

## DOCUMENTS

- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

## GALLERY IMAGES



**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**