
www.vectorlabs.com
Email: customerservice@vectorlabs.com
Telephone: (650) 697-3600

# AZIDO-DPEG®$_{12}$-TFP ESTER

**SKU:** QBD-10569




## DESCRIPTION

Azido-dPEG®12-TFP ester, product number QBD-10569, is a crosslinking compound designed for copper(I)-catalyzed, ruthenium catalyzed, and strain promoted click chemistry (CuAAC, RuAAC, and SPAAC, respectively). The two ends of the molecule are separated by a medium-length (40 atoms), single molecular weight, discrete polyethylene glycol (dPEG®) spacer. The spacer imparts water solubility to QBD-10569 and increases the hydrodynamic volume of the conjugate molecule. The azide group provides the click chemistry functionality, while the 2,3,5,6-tetrafluorophenyl (TFP) ester reacts primary and secondary amines. The single molecular weight and discrete chain length of the spacer simplifies analysis of the product and of its conjugates.

TFP esters are an alternative to the widely popular N-hydroxysuccinimidyl (NHS) esters. TFP esters are more hydrolytically stable than NHS esters, and their optimal reaction pH range (7.5 – 8.0) is somewhat higher than NHS esters (7.0 – 7.5). Moreover, TFP esters in their optimal pH range are more reactive toward free amines than NHS esters.

## SPECIFICATIONS

| | |
|---|---|
| **CAS Number** | N/A |
| **Molecular Weight** | 791.78; single compound |
| **Chemical Formula** | $C_{33}H_{53}F_4N_3O_{14}$ |

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**


www.vectorlabs.com
Email: customerservice@vectorlabs.com
Telephone: (650) 697-3600

| | |
|---|---|
| **Purity** | > 98% |
| **Unit Size** | 100mg, 1000mg |
| **Solubility** | Methylene Chloride, Acetonitrile, Ethyl Acetate, THF, Methanol, Isopropanol, or Water |
| **Spacers** | dPEG® Spacer is 40 atoms and 46.306 Å |
| **Storage Instructions** | -20°C; Always let come to room temperature before opening; be careful to limit exposure to moisture and restore under an inert atmosphere; stock solutions can be prepared with dry solvent and kept for several days (freeze when not in use). dPEG® pegylation compounds are generally hygroscopic and should be treated as such. This will be less noticeable with liquids, but the solids will become tacky and difficult to manipulate, if care is not taken to minimize air exposure. |
| **Shipping Instructions** | Ambient |

# DOCUMENTS

- Safety Data Sheet
- Datasheet

# GALLERY IMAGES




**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**