




BIOTIN-DPEG®₁₁-NH₂

SKU: QBD-10196



DESCRIPTION

Biotin-dPEG®₁₁-NH₂ (Biotin-dPEG®₁₁-amine), product number QBD-10196, is a medium-length, hydrophilic, water-soluble, discrete PEG biotinylation product that reacts with carbonyls, carboxylic acids, and the activated esters (N-hydroxysuccinimide, 2,3,5,6-tetrafluorophenyl) of carboxylic acids. Additionally, Biotin-dPEG®₁₁-NH₂ couples directly to a carboxylic acid using 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) chemistry. Furthermore, the product can react with carbonyl groups (aldehydes or ketones) to form a Schiff base, which can be reduced to a secondary amine using sodium cyanoborohydride if additional conjugate stability is necessary.

Amphiphilic biotin-dPEG®₁₁-NH₂ dissolves equally well in both aqueous buffer and organic solvent. It is not necessary to dissolve this product in an organic solvent before using it in aqueous reaction media. Furthermore, biotinylation of biomacromolecules with QBD-10196 will not cause aggregation and precipitation. Eliminating aggregation should consequently improve the signal-to-noise ratio of applications developed with this product. Reaction with a carboxylic acid or active ester of a carboxylic acid will yield a stable amide bond. Aldehydes are uncommon in biomacromolecules. However, the oxidation of reducing sugars in glycosylated proteins forms aldehydes in glycosylated proteins. Reactions with aldehydes will produce labile imines or Schiff bases that can be reduced under mild conditions with sodium cyanoborohydride to form more stable secondary amines.

Any application that can take advantage of the strong biotin-avidin/ streptavidin affinity can use Biotin-dPEG®₁₁-NH₂. The terminal amine of QBD-10196 offers the opportunity to biotinylate (1) the carbohydrate coat of glycoproteins and (2) surfaces consisting of carboxylic acids or their active esters. Thus, Biotin-dPEG®₁₁-NH₂ expands the range of molecules and nanoparticles for

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



which dPEG®-biotin is useful.


SPECIFICATIONS

CAS Number	604786-74-5
Molecular Weight	770.97; single compound
Chemical Formula	C ₃₄ H ₆₆ N ₄ O ₁₃ S
Purity	> 98%
Unit Size	100 mg, 1000 mg
Solubility	Methylene chloride, DMAC, DMSO or water.
Spacers	dPEG® Spacer is 37 atoms and 44.1 Å
Storage Instructions	-20°C; Always let come to room temperature before opening; be careful to limit exposure to moisture and restore under an inert atmosphere; stock solutions can be prepared with dry solvent and kept for several days (freeze when not in use). dPEG® pegylation compounds are generally hygroscopic and should be treated as such. This will be less noticeable with liquids, but the solids will become tacky and difficult to manipulate, if care is not taken to minimize air exposure.
Shipping Instructions	Ambient

DOCUMENTS

- [Safety Data Sheet](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.