



GOAT ANTI-MOUSE IGG ANTIBODY (H+L), BIOTINYLATED

SKU: BA-9200-1.5



DESCRIPTION

The biotinylated goat anti-mouse IgG secondary antibody is used in an avidin-biotin or streptavidin-biotin detection system. These antibodies can be used for tissue and cell staining, ELISAs, and blots.

Features:

- Recognizes both heavy and light chains (H+L)
- Supplied in solution

SPECIFICATIONS

Format

Concentrate

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



| | |
|-----------------------------|--|
| Formulation | 10 mM sodium phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide, 3 mg/ml bovine serum albumin. |
| Unit Size | 1.5 mg |
| Storage Instructions | 2-8 °C; Store frozen for long term storage |
| Usage Summary | The recommended concentration range for use is 2-10 µg/ml. If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue. NOTE: This product is raised in goat and is not the same as that contained in the VECTASTAIN ABC Kits for the detection of mouse IgG which is raised in horse. |
| Applications | Immunohistochemistry / Immunocytochemistry, Immunofluorescence, In situ hybridization, Blotting Applications, Elispot, ELISAs |
| Target Species | Mouse |
| Concentration | 1.5 mg active conjugate/ml |
| Conjugate | Biotinylated |
| Reactive Species | Goat |
| Source Species | Mouse |
| Host Species | Goat |

TECHNICAL INFORMATION

Vector Laboratories affinity-purified antibodies are of unmatched quality. These antibodies are prepared using proprietary immunization schedules that produce high affinity antibodies. The antibodies are then purified by affinity chromatography, and cross-reactivities that are likely to interfere with specific labeling are removed by solid phase adsorption techniques. The biotinylated secondary antibodies are conjugated to ensure the maximum degree of labeling without compromising the specificity or affinity of the antibody.

With some exceptions the recommended dilution for most applications is 1:200.

CITATIONS

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



Powered by Bioz © 2023 See more details on Bioz

DOCUMENTS

- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.