



GRIFFONIA SIMPLICIFOLIA LECTIN I (GSL I) ISOLECTIN B₄, BIOTINYLATED

SKU: B-1205-.5



DESCRIPTION

GSL I-B₄ isolectin contains only the B subunits. It is a useful marker for endothelial cells from nonprimates such as mouse, rat, rabbit, and goat as well as a marker for non-peptidergic unmyelinated primary afferent neurons. This “B”-rich lectin preferentially agglutinates blood group B cells and is specific for α -galactose residues.

Biotinylated GSL I-B₄ has an appropriate number of biotins bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated biotins and is preserved with sodium azide.

SPECIFICATIONS

Molecular Weight	114
-------------------------	-----

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



Extinction Coefficient	1.4
Formulation	10 mM HEPES, pH 7.5, 0.15 M NaCl, 0.08% sodium azide, 0.1 mM CaCl ₂ .
Inhibiting or Eluting Sugar	Galactose
Unit Size	0.5 mg
Storage Instructions	2-8 °C; Store frozen for long term storage
Sugar Specificity	Terminal αGal
Usage Summary	For most applications, we recommend a freshly prepared working solution of 5-20 µg/ml in the below buffer.
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, Blotting Applications, Elispot, ELISAs, Glycobiology
Concentration	0.5 mg active conjugate/ml
Conjugate	Biotinylated

TECHNICAL INFORMATION

GSL I is a family of glycoproteins with molecular weights of approximately 114 kDa. There are two types of subunits, termed A and B, with slightly different molecular weights. These subunits combine to form tetrameric structures, resulting in five isolectins. The A-rich lectin preferentially agglutinates blood group A erythrocytes and thus appears to be specific for α-N-acetylgalactosamine residues, while the B-rich lectin preferentially agglutinates blood group B cells and is specific for α-galactose residues. Our GSL I is a mixture of the five isolectins. GSL I has been reported to bind several glycoproteins including laminin.

This biotinylated lectin is an ideal intermediate for examining glycoconjugates using the Biotin-Avidin/Streptavidin System. First the biotinylated lectin is added, followed by the VECTASTAIN ABC Reagent, Avidin D conjugate, or streptavidin derivative.

Inhibiting Sugar: 500 mM galactose or 100 mM raffinose

CITATIONS



Powered by Bioz © 2023 See more details on Bioz

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



DOCUMENTS

- [Safety Data Sheet](#)
- [Lectins in Histochemistry, ELISA, and Western Blot Applications](#)
- [Download CoA](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.