


BROMOACETAMIDO-DPEG®₄-TFP ESTER

SKU: QBD-11200



Bromoacetamido-dPEG®₄-TFP ester, product number QBD-11200, is a short (18 atoms, 20.4 Å), single molecular weight, discrete PEG (dPEG®) crosslinker functionalized with a thiol-reactive bromoacetate group on one end and an amine-reactive 2,3,5,6-tetrafluorophenyl (TFP) ester on the other.

The short, flexible, non-immunogenic dPEG® crosslinker increases the hydrodynamic volumes of conjugates in aqueous media, thereby increasing water solubility. The increased hydrodynamic volume can reduce or eliminate conjugate molecules' renal clearance and shield conjugates from opsonization.

The bromoacetate moiety is an alternative to the maleimide group, reacting chemoselectively with free thiols at pH ≥ 8.0 . It is useful for situations where the conjugate molecule needs to be kept at a high pH (≥ 8.0), above the range where maleimides can be used effectively, or in constructs where maleimide groups are unstable.

TFP esters are an alternate reactive group to N-hydroxysuccinimidyl (NHS) esters and react with amines at an optimal pH range of 7.5 - 8.0, higher than the optimal range for NHS esters (pH 7.0 - 7.5). TFP esters are more hydrolytically stable than NHS esters, and they are more reactive toward amines in their optimal pH range than NHS esters are within the pH range 7.0 - 7.5.

Specifications

Unit Size	100 mg, 1000 mg
Molecular Weight	534.29; single compound;
Chemical formula	C ₁₉ H ₂₄ BrF ₄ NO ₇
CAS	N/A

For research use only. Not intended for animal or human therapeutic or diagnostic use.

Purity	> 97%
Spacers	dPEG® Spacer is 18 atoms and 20.4 Å
Shipping	Ambient
Typical solubility properties (for additional information contact Customer Support)	Methylene Chloride, Methanol, MTBE, or Water.
Storage and handling	-20°C; Always let come to room temperature before opening; be careful to limit exposure to moisture and restore under an inert atmosphere; stock solutions can be prepared with dry solvent and kept for several days (freeze when not in use). dPEG® pegylation compounds are generally hygroscopic and should be treated as such. This will be less noticeable with liquids, but the solids will become tacky and difficult to manipulate, if care is not taken to minimize air exposure.

For research use only. Not intended for animal or human therapeutic or diagnostic use.