




## **GRIFFONIA SIMPLICIFOLIA LECTIN I (GSL I) ISOLECTIN B<sub>4</sub>, DYLIGHT 594**

**SKU:** DL-1207-.5



---

### **DESCRIPTION**

GSL I-B<sub>4</sub> isolectin contains only the B subunits. It is a useful marker for endothelial cells from nonprimates such as mouse, rat, rabbit, and goat as well as a marker for non-peptidergic unmyelinated primary afferent neurons. This “B”-rich lectin preferentially agglutinates blood group B cells and is specific for  $\alpha$ -galactose residues.

DyLight™ 594 GSL I-B<sub>4</sub> isolectin is produced by using special conjugation procedures to incorporate DyLight™ 594 into our affinity-purified lectin. This conjugate has an appropriate number of fluorochromes bound which provide the maximum fluorescence and optimum staining characteristics for this particular lectin. This reagent is supplied essentially free of unconjugated fluorochromes and inactive lectin.

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



## SPECIFICATIONS

<b>Molecular Weight</b>	114
<b>Color of Fluorescence</b>	Red
<b>Extinction Coefficient</b>	1.4
<b>Formulation</b>	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM CaCl <sub>2</sub> .
<b>Inhibiting or Eluting Sugar</b>	Galactose
<b>Maximum Emission</b>	617 nm
<b>Maximum Excitation</b>	592 nm
<b>Unit Size</b>	0.5 mg
<b>Storage Instructions</b>	2-8 °C
<b>Sugar Specificity</b>	Terminal $\alpha$ Gal
<b>Usage Summary</b>	The recommended concentration range for use is 5-20 $\mu$ g/ml.
<b>Applications</b>	Immunofluorescence, Glycobiology
<b>Concentration</b>	1 mg active conjugate/ml
<b>Conjugate</b>	DyLight 594

## TECHNICAL INFORMATION

GSL I is a family of glycoproteins with molecular weights of approximately 114 kDa. There are two types of subunits, termed A and B, with slightly different molecular weights. These subunits combine to form tetrameric structures, resulting in five isolectins. The A -rich lectin preferentially agglutinates blood group A erythrocytes and thus appears to be specific for  $\alpha$ -N-acetylgalactosamine residues, while the B -rich lectin preferentially agglutinates blood group B cells and is specific for  $\alpha$ -galactose residues. Our GSL I is a mixture of the five isolectins. GSL I has been reported to bind several glycoproteins including laminin.

Accompanying each fluorescent lectin is an analysis data sheet summarizing the results of our quality control tests and providing pertinent information on the product. All of these reagents are supplied as solutions preserved with sodium azide.

- Excitation maximum: 592 nm
- emission maximum: 617 nm
- Color: Red

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



Inhibiting Sugar: 500 mM galactose or 100 mM raffinose

## CITATIONS




Powered by Bioz © 2023 See more details on Bioz

## DOCUMENTS

- [Lectins in Histochemistry, ELISA, and Western Blot Applications](#)
- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

## GALLERY IMAGES



**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**