



# **GRIFFONIA SIMPLICIFOLIA LECTIN I (GSL I) ISOLECTIN B4, DYLIGHT 649**

**SKU:** DL-1208-.5



---

## **DESCRIPTION**

GSL I-B<sub>4</sub> isolectin contains only the B subunits. It is a useful marker for endothelial cells from nonprimates such as mouse, rat, rabbit, and goat as well as a marker for non-peptidergic unmyelinated primary afferent neurons. This “B”-rich lectin preferentially agglutinates blood group B cells and is specific for  $\alpha$ -galactose residues.

DyLight™ 649 GSL I-B<sub>4</sub> isolectin is produced by using special conjugation procedures to incorporate DyLight™ 649 into our affinity-purified lectin. This conjugate has an appropriate number of fluorochromes bound which provide the maximum fluorescence and optimum staining characteristics for this particular lectin. This reagent is supplied essentially free of unconjugated fluorochromes and inactive lectin.

- Excitation maximum: 646 nm
- emission maximum: 674 nm

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



- Color: Red

## SPECIFICATIONS

<b>Molecular Weight</b>	114
<b>Color of Fluorescence</b>	Far Red
<b>Extinction Coefficient</b>	1.4
<b>Formulation</b>	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM CaCl <sub>2</sub> .
<b>Inhibiting or Eluting Sugar</b>	Galactose
<b>Maximum Emission</b>	670 nm
<b>Maximum Excitation</b>	655 nm
<b>Unit Size</b>	0.5 mg
<b>Storage Instructions</b>	2-8 °C
<b>Sugar Specificity</b>	Terminal $\alpha$ Gal
<b>Usage Summary</b>	The recommended concentration range for use is 5-20 $\mu$ g/ml. If a precipitate forms upon long-term storage, warm to 37 °C.
<b>Applications</b>	Immunofluorescence, Glycobiology
<b>Concentration</b>	1 mg active conjugate/ml
<b>Conjugate</b>	DyLight 649

## TECHNICAL INFORMATION

GSL I is a family of glycoproteins with molecular weights of approximately 114 kDa. There are two types of subunits, termed A and B, with slightly different molecular weights. These subunits combine to form tetrameric structures, resulting in five isolectins. The A -rich lectin preferentially agglutinates blood group A erythrocytes and thus appears to be specific for  $\alpha$ -N-acetylgalactosamine residues, while the B -rich lectin preferentially agglutinates blood group B cells and is specific for  $\alpha$ -galactose residues. Our GSL I is a mixture of the five isolectins. GSL I has been reported to bind several glycoproteins including laminin.

Accompanying each fluorescent lectin is an analysis data sheet summarizing the results of our quality control tests and providing pertinent information on the product. All of these reagents are supplied as solutions preserved with sodium azide.

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



Inhibiting Sugar: 500 mM galactose or 100 mM raffinose

## DOCUMENTS

- [Safety Data Sheet](#)
- [Lectins in Histochemistry, ELISA, and Western Blot Applications](#)
- [Download CoA](#)
- [Datasheet](#)

## GALLERY IMAGES



**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**