



GOAT ANTI-RABBIT IGG ANTIBODY (H+L), FLUORESCEIN

SKU: FI-1000-1.5



DESCRIPTION

Fluorescein Goat Anti-Rabbit IgG Antibody can be used for immunofluorescence and other applications. Optimal F/P ratios have been established for each conjugate to ensure maximum fluorescence with minimal background staining.

Features:

- Recognizes both heavy and light chains (H+L)
- Optimally labeled with fluorescein to provide the brightest label for fluorescence microscopy
- Supplied in solution
- Excitation: 495 nm
- Emission: 515 nm
- Color: Green

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



SPECIFICATIONS

Color of Fluorescence	Green
Format	Concentrate
Formulation	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide.
Maximum Emission	510-520 nm
Maximum Excitation	490-500 nm
Unit Size	1.5 mg
Storage Instructions	2-8 °C
Usage Summary	The recommended concentration range for use is 5-20 µg/ml. If this fluorescein-labeled antibody is to be used in tissues which may contain cross-reacting endogenous immunoglobulins, dilution of this antibody may be made in buffers containing 2% normal serum from the same species as the tissue.
Applications	Immunofluorescence, In situ hybridization, Blotting Applications, Flow Cytometry/Cell Separation
Target Species	Rabbit
Concentration	1.5 mg active conjugate/ml
Conjugate	Fluorescein
Reactive Species	Goat
Source Species	Rabbit
Host Species	Goat

TECHNICAL INFORMATION

The goat anti-rabbit Ig antibodies are prepared by hyperimmunizing animals in a manner that produces high affinity antibodies. These are then purified by an affinity chromatography procedure designed to remove any low affinity antibodies which may be present. Cross-reactivities that are likely to interfere with specific labeling are removed by solid-phase adsorption techniques. The final product is then subjected to rigorous quality control assays including immunodiffusion, solid-phase enzyme immunoassays, gel electrophoresis and solid-phase binding assays. In preparing the labeled antibodies, great care is taken to ensure the maximum degree of labeling with no alteration in the specificity and affinity of the antibody. The labeled antibody then undergoes a further series of quality control assays, including immunohistochemical analysis.

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



Fluorescein Goat Anti-Rabbit IgG (H+L) Antibody can be used for flow cytometry or for tissue staining.

CITATIONS



Powered by Bioz © 2023 See more details on Bioz

DOCUMENTS

- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.