



LENS CULINARIS AGGLUTININ (LCA), FLUORESCIN

SKU: FL-1041-5



DESCRIPTION

LCA recognizes sequences containing α -linked mannose residues but recognizes additional sugars as part of the receptor structure, giving it a narrower specificity than Con A. An α -linked fucose residue attached to the N-acetylchitobiose portion of the core oligosaccharide significantly enhances affinity. By exploiting this narrower specificity, glycoproteins and glycopeptides can be subfractionated with LCA after initial isolation with Con A.

Fluorescein labeled LCA has an appropriate number of fluorochromes bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated fluorochromes. The excitation maximum is at 495 nm and the emission maximum is at 515 nm.

SPECIFICATIONS

Molecular Weight	50
-------------------------	----

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



Color of Fluorescence	Green
Extinction Coefficient	1.25
Formulation	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM CaCl ₂ , 0.01 mM MnCl ₂ .
Inhibiting or Eluting Sugar	Mixture of α -methyl-mannoside & α -methyl-glucoside
Maximum Emission	514-521 nm
Maximum Excitation	495-500 nm
Unit Size	5 mg
Storage Instructions	2-8 °C
Sugar Specificity	α 1,6-linked fucose
Usage Summary	The recommended concentration range for use is 5-20 μ g/ml.
Applications	Immunofluorescence, Glycobiology
Concentration	5 mg active conjugate/ml
Conjugate	Fluorescein

TECHNICAL INFORMATION

Lens culinaris agglutinin is composed of four subunits – two of about 17 kDa and two of 8 kDa. LCA has been found to be one of the most effective agents in preventing skin allograft rejection in model systems. LCA has been used to purify numerous glycoproteins, including immunoglobulins, histocompatibility antigens, and α 2-macroglobulin.

Accompanying each fluorescent lectin is an analysis data sheet summarizing the results of our quality control tests and providing pertinent information on the product. All of these reagents are supplied as solutions preserved with sodium azide.

Inhibiting/Eluting Sugar: mixture of 200 mM α -methylmannoside/200 mM α -methylglucoside

CITATIONS



Powered by Bioz © 2023 See more details on Bioz

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



DOCUMENTS

- [Lectins in Histochemistry, ELISA, and Western Blot Applications](#)
- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.