



# **PHASEOLUS VULGARIS LEUCOAGGLUTININ (PHA-L), FLUORESCIN**

**SKU:** FL-1111-2



---

## **DESCRIPTION**

*Phaseolus vulgaris* agglutinin is the name ascribed to a family of lectins, each of which consists of four subunits. There are two different types of subunits. One appears to be involved primarily in red cell agglutination and has been designated the “E” subunit (for erythroagglutinin). The other type is involved in lymphocyte agglutination and mitogenic activity and has been termed the “L” subunit (for leucoagglutinin). These subunits combine to produce five isolectins. PHA-L, with four “L” type subunits, does not agglutinate red cell but is a potent mitogen.

Fluorescein labeled PHA-L has an appropriate number of fluorochromes bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated fluorochromes. The excitation maximum is at 495 nm and the emission maximum is at 515 nm.

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



## SPECIFICATIONS

<b>Molecular Weight</b>	126
<b>Color of Fluorescence</b>	Green
<b>Extinction Coefficient</b>	1.16
<b>Formulation</b>	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM CaCl <sub>2</sub>
<b>Inhibiting or Eluting Sugar</b>	GalNAc
<b>Maximum Emission</b>	514-521 nm
<b>Maximum Excitation</b>	495-500 nm
<b>Unit Size</b>	2 mg
<b>Storage Instructions</b>	2-8°C
<b>Sugar Specificity</b>	β1,6-branched N-glycan
<b>Usage Summary</b>	The recommended concentration range for use is 5-20 μg/ml.
<b>Applications</b>	Immunofluorescence, Glycobiology
<b>Concentration</b>	2 mg active conjugate/ml
<b>Conjugate</b>	Fluorescein

## TECHNICAL INFORMATION

*PHA-L, with four L type subunits, does not agglutinate red cells but is a potent mitogen. The other three isolectins, designated E3L1, E2L2, and E1L3, have erythroagglutinating and mitogenic activities proportional to the number of respective E or L subunits. We have termed the mixture of the five isolectins PHA (E+L).*

Accompanying each fluorescent lectin is an analysis data sheet summarizing the results of our quality control tests and providing pertinent information on the product. All of these reagents are supplied as solutions preserved with sodium azide.

Elution: 100 mM acetic acid

## CITATIONS

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



Powered by Bioz © 2023 See more details on Bioz

## DOCUMENTS

- [Lectins in Histochemistry, ELISA, and Western Blot Applications](#)
- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

## GALLERY IMAGES



**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**