



SOYBEAN AGGLUTININ (SBA), FLUORESCEIN

SKU: FL-1011-2



DESCRIPTION

SBA preferentially binds to oligosaccharide structures with terminal α - or β -linked *N*-acetylgalactosamine, and to a lesser extent, galactose residues. Binding can be blocked by substitutions on penultimate sugars, such as fucose attached to the penultimate galactose in blood group B substance.

Fluorescein labeled Soybean agglutinin has an appropriate number of fluorochromes bound to provide the optimum staining characteristics for this lectin. This conjugate is supplied essentially free of unconjugated fluorochromes. The excitation maximum is at 495 nm and the emission maximum is at 515 nm.

SPECIFICATIONS

Molecular Weight	120
Color of Fluorescence	Green

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



Extinction Coefficient	1.33
Formulation	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM CaCl ₂
Inhibiting or Eluting Sugar	GalNAc
Maximum Emission	514-521 nm
Maximum Excitation	495-500 nm
Unit Size	2 mg
Storage Instructions	2-8 °C
Sugar Specificity	Terminal GalNAc epitopes and Forssman antigen
Usage Summary	The recommended concentration range for use is 5-20 µg/ml.
Applications	Immunofluorescence, Glycobiology
Concentration	2 mg active conjugate/ml
Conjugate	Fluorescein

TECHNICAL INFORMATION

Composed of four subunits of approximately equal size, soybean agglutinin is a family of closely related isolectins. This glycoprotein has a molecular weight of about 120 kDa and an isoelectric point near pH 6.0.

An important application for SBA is the separation of pluripotent stem cells from human bone marrow. Cells fractionated by SBA do not produce graft vs host disease and can be used in bone marrow transplantation across histocompatibility barriers.

Accompanying each fluorescent lectin is an analysis data sheet summarizing the results of our quality control tests and providing pertinent information on the product. All of these reagents are supplied as solutions preserved with sodium azide.

Inhibiting/Eluting Sugar: 200 mM *N*-acetylgalactosamine

CITATIONS



Powered by Bioz © 2023 See more details on Bioz

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



DOCUMENTS

- [Lectins in Histochemistry, ELISA, and Western Blot Applications](#)
- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.