




HYDROXY-DPEG®₁₂-T-BUTYL ESTER

SKU: QBD-10719



DESCRIPTION

Hydroxy-dPEG®₁₂-t-butyl ester, product number QBD-10719, is a versatile building block based upon a monodispersed polyethylene glycol (PEG) chain. A primary alcohol group terminates one end of the molecule. The opposite end is a propionic acid group protected as the tert-butyl ester. The terminal hydroxy group can be functionalized with various reactive groups to permit direct modification of surfaces, small molecules, or biomolecules. Alternatively, the hydroxy group can react with isothiocyanates or active esters of carboxylic acids to form new conjugates. The tert-butyl ester deprotects with trifluoroacetic acid (TFA) or formic acid, exposing the terminal propionic acid, which can then be activated as the NHS or TFP ester or directly coupled to an amine using a suitable carbodiimide.

SPECIFICATIONS

CAS Number	1186025-29-5
Molecular Weight	674.81; single compound
Chemical Formula	C ₃₁ H ₆₂ O ₁₅
Purity	> 98%
Unit Size	100mg, 1000mg
Solubility	Methylene chloride, Acetonitrile, DMAC or DMSO.
Spacers	dPEG® Spacer is 40 atoms and 46.4 Å

For research use only. Not intended for therapeutic or diagnostic use in animals or humans.



Storage Instructions


-20°C; Always let come to room temperature before opening; be careful to limit exposure to moisture and restore under an inert atmosphere; stock solutions can be prepared with dry solvent and kept for several days (freeze when not in use). dPEG® pegylation compounds are generally hygroscopic and should be treated as such. This will be less noticeable with liquids, but the solids will become tacky and difficult to manipulate, if care is not taken to minimize air exposure.

Shipping Instructions Ambient

DOCUMENTS

- [Safety Data Sheet](#)
- [Datasheet](#)

GALLERY IMAGES



For research use only. Not intended for therapeutic or diagnostic use in animals or humans.