



# GOAT ANTI-RABBIT IGG ANTIBODY (H+L), TEXAS RED

**SKU:** TI-1000-1.5



---

## DESCRIPTION

The Texas Red Goat Anti-Rabbit IgG Antibody can be used for immunofluorescence and other applications. Optimal F/P ratios have been established for each conjugate to ensure maximum fluorescence with minimal background staining.

### Features:

- Recognizes both heavy and light chains (H+L)
- Optimally labeled with Texas Red to provide the brightest label for fluorescence microscopy
- Supplied in solution
- Excitation: 595 nm
- Emission: 615 nm
- Color: Red

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



## SPECIFICATIONS

**Color of Fluorescence** Red

**Format** Concentrate

**Formulation** 10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide

**Maximum Emission** 606-615 nm

**Maximum Excitation** 595-604 nm

**Unit Size** 1.5 mg

**Storage Instructions** 2-8 °C

**Usage Summary** The recommended concentration range for use is 5-20 µg/ml. If this antibody is to be used in tissues which may contain cross-reacting endogenous immunoglobulins, dilution of this antibody may be made in buffers containing 2 % normal serum from the same species as the tissue.

**Applications** Immunofluorescence, In situ hybridization, Blotting Applications, Flow Cytometry/Cell Separation

**Target Species** Rabbit

**Concentration** 1.5 mg active conjugate/ml

**Conjugate** Texas Red

**Reactive Species** Goat

**Source Species** Rabbit

**Host Species** Goat

## TECHNICAL INFORMATION

The goat anti-rabbit Ig antibodies are prepared by hyperimmunizing animals in a manner that produces high affinity antibodies. These are then purified by an affinity chromatography procedure designed to remove any low affinity antibodies which may be present. Cross-reactivities that are likely to interfere with specific labeling are removed by solid-phase adsorption techniques. The final product is then subjected to rigorous quality control assays including immunodiffusion, solid-phase enzyme immunoassays, gel electrophoresis and solid-phase binding assays. In preparing the labeled antibodies, great care is taken to ensure the maximum degree of labeling with no alteration in the specificity and affinity of the antibody. The labeled antibody then undergoes a further series of quality control assays, including immunohistochemical analysis.

**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**



Texas Red (sulforhodamine 101) excites at about 595 nm and emits at about 615 nm, producing a bright red fluorescence well removed from the emission spectrum of fluorescein. Texas Red Goat Anti-Rabbit IgG (H+L) Antibody conjugate is ideal as a single or double label reagent for flow cytometry or tissue staining.

## CITATIONS



Powered by Bioz © 2023 See more details on Bioz

## DOCUMENTS

- [Safety Data Sheet](#)
- [Download CoA](#)
- [Datasheet](#)

## GALLERY IMAGES



**For research use only. Not intended for therapeutic or diagnostic use in animals or humans.**